Measurements and Analysis of the Thermal Conductivity Close to the Vapor-Liquid Critical Line of Binary Fluid Mixtures

E.P. Sakonidou, H.R. van den Berg, and C.A. ten Seldam

Van der Waals-Zeeman Institute

University of Amsterdam

Valckenierstraat 67

1018 XE Amsterdam, The Netherlands

J.V. Sengers

Department of Chemical Engineering and Institute for Physical Science and Technology

University of Maryland

College Park, MD 20742 USA

It has been well established theoretically and experimentally that the thermal conductivity of one-component fluids shows a strong enhancement close to the vapor-liquid critical point and ultimately becomes infinite [1,2]. In case of binary fluid mixtures, already since 1971, several theories predict that asymptotically close to the line of vapor-liquid critical points, also referred to as plait points, the critical enhancement of the thermal conductivity remains finite [3]. However, the available thermal-conductivity data on binary fluid mixtures did not offer any experimental evidence for the predicted asymptotic behavior [4]. The most recent measurements were made on carbon dioxide-ethane mixtures with the guarded parallel-plate apparatus [5]. But also these measurements showed a behavior comparable to that found for one-component fluids and, therefore, failed to demonstrate the occurrence of the finite thermal conductivity near the plait point.

We present now new experimental thermal-conductivity data of an equimolar methane-ethane mixture very close to the plait point. For these measurements the upgraded version of our parallel-plate apparatus [6] was used. Since these new data on the equimolar mixture showed a small uncertainty, even in the near vicinity of the plait point, they allowed us to confirm for the first time experimentally that the thermal conductivity of a mixture does not diverge but crosses over to a finite asymptotic behavior at the plait point. These data were analyzed in terms of a new extension of the mode coupling theory for the transport properties, which includes the non-asymptotic critical behavior [7]. As result of this analysis, excellent agreement was found between measured and calculated thermal-conductivity values. A separate analysis was made on earlier measurements for methane-ethane mixtures, obtained with a different method [4], for various compositions and relatively far from the plait point. This analysis revealed the same qualitative behavior for the thermal conductivity, however, quantitatively large differences were found.

- [1] J. Luettmer-Strathmann, J.V. Sengers, and G.A. Olchowy, J. Chem. Phys. 103, 7482 (1995).
- [2] E.P. Sakonidou, H.R. van den Berg, C.A. ten Seldam ,and J.V. Sengers, J. Chem. Phys. 105, 10535 (1996).
- [3] M.A. Anisimov, A.V. Voronel, and E.E. Gorodestkii, Sov. Phys. JETP 33, 605 (1971).
- [4] H.M. Roder and D. G. Friend, *Int. J. Thermophys.* **6**, 607 (1985); *Experimental Thermal-Conductivity Values for Mixtures of Methane and Ethane*, NBSIR 85-3024 (National Bureau of Standards (U. S.) Interagency Report 1985).
- [5] R. Mostert, H.R. van den Berg, P. S. van der Gulik, and J. V. Sengers, High Temp.-High Press. 24, 469 (1992).
- [6] E. Sakonidou, Ph.D. Thesis (Van der Waals-Zeeman Institute, Univ. of Amsterdam, 1996).
- [7] J. Luettmer-Strathmann and J. V. Sengers, J. Chem. Phys. 104, 3026 (1996).